

WDOWIAK, Szymon

The recent-carrie glacier in the Great Mieguszwiecki Kettle
over Morskie Oko Village in the Tatra Mountains. Biuletyn Geolog
1 no.1:87-92 '61.

1. Chair of Quaternary Geology, University, Warsaw.

WDOWIAK, T.

Optimal conditions of heating bales of greasy wool. Biuletyn Wlok. p.9.

(PRZEMYSŁ WLOKIENNICZY. Vol. 11, No. 5, May 1957. Warszawa, Poland)

SO: Monthly List of East European Accessions (EEAL) LC. Vol. 6, No. 10, October 1957. Uncl.

WDOWIAK, Tadeusz; PIATKOWSKA, Teresa

New positioning of the fancy roller. Przegl włokien
16 no.9:468-471 S '62.

WDOWIAK, Tadeusz, mgr. inż.

"New textile technologies, nonwoven fabrics." Reviewed
by Tadeusz Wdowiak. Przegl włokien 17 no. 2:102-103 P '63.

KAMIENSKA, Karolina; WDOWIAK, Wanda; KRAJEWSKI, Stanislaw

A case of Bleck-Sulzberger syndrome. Pediat pol 36 no.10:1079-1086
O '61.

1. Z II Kliniki Poloznictwa i Chorob Kobietych AM w Gdansk Kierownik:
doc. dr med. W Gremadzki i z Oddzialu Dermatologii Szpitala Wojewodzkiego
im. M. Kopernika w Gdansk Ordynator: dr med. S.Krajewski.
(PIGMENTATION DISORDERS in inf & child)

KAMIENSKA, Karolina; WDOWIAK, Wania

Acute toxic methemoglobinemia in newborn infants. Ginek. pol. 33 no.5:
617-622 '62.

1. Z II Kliniki Położnictwa i Chorob Kobietych AM w Gdańsku. Kierownik:
doc. dr med. W. Gromadzki.
(METHEMOGLOBINEMIA) (INFANT NEWBORN DISEASES)

HEJKA, Zuzanna; JANIKOWSKI, Tadeusz; KRYWKO, Alina; TYLICKA, Teresa;
WDOWIAK, Wanda; WOZNICZKO, Jerzy.

Incidence of neurologic symptoms in the newborns in relation
to causative factors. (Ginek. Pol. 36 no. 12:1379-1386 D ' 65

1. Z Kliniki Neurologicznej AM w Gdansk (Kierownik: prof. dr.
med. Z. Majewska); z I Kliniki Poloznictwa i Chorob Kobietych
AM w Gdansk (Kierownik: prof. dr. med. S. Metler) i z II
Kliniki Poloznictwa i Chorob Kobietych AM w Gdansk (Kierownik:
prof. dr. med. W. Gromadzki).

WDOWIANZ, J.

"Geologic Exploration for Petroleum Deposits", (To be contd.) p. 163, (NAPTA,
Vol. 3, No. 6, June 1952, Krakow, Poland)

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, No. 5,
May 1955, Uncl.

WIDOMIARZ, J.

"Geologic Exploration for Petroleum Deposits", (To be contd.) p. 194,
(NAPTA, Vol. 8, No. 7, July 1952, Krakow, Poland)

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, No. 5,
May 1955, Uncl.

Wdowiarsz, s.

Crude oil and natural gas in relation to the geology of the Carpathian Mountains:
a summary of a report.

P. 456 (Przegląd Geologiczny, Vol. 4, no. 10, Oct. 1956, Warszawa, Poland)

Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 2,
February 1958

WDOWIARZ, S.

Problems of oil in northern and central Poland. p. 541.
(PRZEGLAD GEOLOGICZNY. Vol. 4, no. 12, Dec. 1956, Warszawa, Poland)

SO: Monthly List of East European Accessions (EEAL) LC. Vol. 6, No. 12, Dec. 1957.
Uncl.

WDOWIARZ, Stanislaw

Jaslo shales in the eastern Carpathian Mountains of Rumania.
Kwartalnik geol 3 no.3:563-568 '59. (EEAI 9:7)

1. Karpacka Stacja Terenowa Instytut Geologicznego
(Carpathian Mountains)
(Rum--Shale)

BOGDANOV, A.A. (SSSR); SLAVIN, V.I. (SSSR); KSIAZKIEWICZ, M. (Pologne);
VARENTSOV, M.I. (SSSR); WIDOWIARZ, St. (Polska); PASHCHENKO, Ya.Ye.
(SSSR); MISHUNINA, Z.A. (SSSR); ZIELINSKI, J. (Polen)

Participation in discussions. Mat.Karp.-Balk.assots. no.1:190-207
'60. (MIRA 14:12)

(Geology)

VYALOV, O.S. (SSSR); MASLOV, V.P. (SSSR); WDOWIARZ, St. (Polska);
OLEWICZ, Z.R. (Polska); NOVAK, V. (Pol'sha); ~~SLAVIN~~, V.I. (SSSR)
MASIAKOVA, N.I. (SSSR); VYALOV, O.S. (SSSR); EBERZIN, A.G. (SSSR)
BONDARCHUK, V.G. (SSSR)

Participation in discussions. Mat.Karp.-Balk.assots. no.3:157-
179 '60. (MIRA 14:12)

(Carpathian Mountains → Geology)

WDOWIARZ, Stanislaw

Oil and natural gas deposits on the territory of the Carpathian Mountains and perspectives for prospecting on that territory. Przegl geol 8 no.10:518 0 '60. (EEAI 10:9)

1. Akademia Gorniczo-Hutnicza.

(Poland---Petroleum) (Poland---Gas, Natural)

WDOWIARZ, S.

Wojew. Przegląd Geologiczny. Vol 10, No 2 (1977). February 1981 (continued)

16

11. "Wzrost i rozwój górnictwa w Polsce" [Growth and Development of Mining in Poland] pp 101-103.
12. "Geologia i geografia województwa łódzkiego" [Geology and Geography of the Łódź Voivodeship] as the University of Łódź, Jan 1978, pp 103-104.
13. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 105-106. (English summary).
14. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 107-108. (English summary).
15. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 109-110.
16. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 111-112.
17. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 113-114.
18. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 115-116.
19. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 117-118.
20. "Pracownicy nauki i sztuki województwa łódzkiego" [Scientists and Artists of the Łódź Voivodeship] pp 119-120.

1197

s/a

WDOWIARZ, Stanislaw

5th Congress of the Carpatho-Balkanian Geological Association. Przegl geolog
10 no. 2:111-118 F. '62.

1. Instytut Geologiczny, Warszawa.

WDOWIARZ, Stanislaw

Geological structure of the Flysch zone of the Eastern Carpathian Mountains and its relationship to the Polish Carpathians.
Kwartalnik geol 5 no.4:991-992 '61. .

1. Karpacka Stacja Terenowa, Instytut Geologiczny, Warszawa.

WDOWIARZ, Stanislaw

POLAND

WDOWIARZ, Stanislaw

Carpathian Field Station, Geological Institute
(Karpacka Stacja Terenowa Instytutu Geologicznego)

Warsaw, Kwartalnik geologiczny, No 3, 1963, pp 542-45.

"Geology of Yugoslavia".

WDOWIARZ, Stanislaw

Report on the 34th Congress of the Polish Geologic Society.
Rocz geol Krakow 32 no.4:681-683 '62.

WDOWIK, M.

2885

Wdowik M. Development of a Closed System of Paper Machine White Water.

176,605

Realizacja zamkniętego obiegu wód maszyny papierniczej. Przegląd Papierniczy. No. 8, 1953, pp. 244-248, 5 figs.

In order to improve operating results, to economise fresh water, and to reduce the consumption of raw materials, a closed whitewater system was evolved and then applied to a board machine. A special purifying device was necessary, and the whitewater returned to the production process was heated up to 40°C in a heat exchanger. Since water containing pulp fragments was used for the sprays washing the wire, a special non-clogging spray pipe, invented by Wdowik, was applied. The closed whitewater system also provides for disinfection by means of a 16% sodium hypochloride which inhibits the development of bacteria.

WDOWINSKI, ZDZISLAŹ.

Amidst forests and lakes. (Translator: Maria Paczynska, photos:
Zdzisław Wdowski) Warszawa, "Sport i Turystyka", 1955.
166 p. (In English. Tr. from the Polish. illus.) SCIENCE

SO: MONTHLY LIST OF EAST EUROPEAN ACCESSIONS (EAL), LC, Vol. 5, no. 2
Feb. 1956

PAVLICEK, Z.; WDOWKA, K.

Traumatic hemobilia. Rozhl. chir. 41 no.2:135-138 P '62.

1. Chirurgické oddelení OUNZ v Prostějově, přednosta MUDr. Mil. Hel.

(BILIARY TRACT wds & inj) (BILE)

HUNGARY

FAZEKAS, Arpad, G., WEBB, James, I., SYMINGTON, Thomas; Medical University of Szeged, Institute of Biochemistry (Szegedi Orvostudományi Egyetem, Biokémiai Intézet), and Pathological Institute of the University, Steroid Research Laboratory, Royal Infirmary, Glasgow.

"In Vitro Study of the Biosynthesis of Corticosteroids in Conn Adenoma Tissue by Using Labelled Precursors."

Budapest, Kiserletes Orvostudomány, Vol XVIII, No 5, Oct 66, pages 480-487.

Abstract: [Authors' Hungarian summary] The biosynthesis of aldosterone and other corticosteroids was studied in vitro by the incubation with labelled steroid precursors of an adrenocortical-adenoma tissue which was the cause of Conn's syndrome in the patient. The steroids produced were identified by radiochemical methods after their isolation by means of paper chromatography. According to the results, a large amount of aldosterone was produced in the adenoma tissue from corticosterone via an 18-OH-corticosterone intermediary. The tritium-labelled 11-dehydrocorticosterone was converted into aldosterone, indicating the role of this steroid as a precursor. Finally, the ratio of corticosterone-cortisol synthesis was shifted in favor of the corticosterone. According to the above results, the cells of the Conn type adenoma show hybrid properties with respect to their biosynthetic capacity since they can synthesize cortisol as well as aldosterone. 1 Hungarian, 24 Western references.
1/1 [Manuscript received 6 Oct 65.]

- 6 -

SVORAD, D., WEBB, W.B., SMIESKY, J.; Pathological Physiology, Slovak Academy of Sciences (Normálne a Patologické Fyziologie SAV), Bratislava.

"Differences in the EEG Sleep Rhythms of Man and Rat."

Prague, Ceskoslovenska Fysiologie, Vol 15, No 2, Feb 66, pp 133-134

Abstract: 16 healthy men, 21 to 31 years old, and 8 rats were investigated with EEG for 3 successive nights. In man, the length of individual stages of sleep is permanent, in rat it changes. The sequence of the stages of sleep and their distribution in certain time intervals are very similar in man and rat. 1 Figure, 2 Western references. Submitted at the "4th Czechoslovak Bioclimatical Conference" at Brno, 11 Oct 65.

1/1

- 127 -

HALMAGYI, D.,; FELKAI, B.,; SOVENYI, E.,; WEBER, A.,; CZIPOTT, Z.,;
KOVACS, G.,; STEINER, B.

Pulmonary circulation in essential pulmonary hemosiderosis. Magyar.
belorv. arch. 8 no.6:188-191 Dec. 55

1. A szegedi Orvostudományi Egyetem I. Belklinikája (igaz: Hetenyi
Geza dr. akadémikus) Szívbetegosztályának közl.

(HEMOCHROMATOSIS

hemosiderosis, essential pulm., pulm. circ. in,
relation to hypertension (Hun))

(BLOOD CIRCULATION

pulm., in essential pulm. hemosiderosis (Hun))

(HYPERTENSION

pulm., relation to essential pulm. Hemosiderosis (Hun))

MIHAI, N., dr.; WEBER, Agathe, asist.

Research on acute and chronic carbon monoxide poisoning.
Changes in arterial pressure. Med. intern. (Bucur) 16
no.9:1113-1119 S '64.

1. Lucrare efectuata in Policlinica combinatului in Centrul
de medicina a muncii, Hunedocara.

WEBER, Apollonia, dr.; BARTOK, Istvan, dr.

A case of essential pulmonary hemosiderosis in adult. Orv.
hetil. 98 no.16:413-418 21 Apr 57.

1. A Szegedi Orvostudományi Egyetem I. sz. Belklinikájának
(igazgató: Hetenyi, Géza, dr. egyet. tanár) és Kórházának és
Kórházvezetői Intézetének (igazgató: Kórpássy, Béla, dr. egyet.
tanár) közleménye.

(HEMOSIDEROSIS, case reports
pulm., essential (Han))

(LUNG DISEASES, case reports
hemosiderosis, essential (Han))

9 (9)

PHASE I BOOK EXPLOITATION

Czech/1562

Weber, Antonín

Velmi krátké vlny (Microwaves) Prague, SNTL, 1957. 335 p.
(Series: Malá elektrotechnická knihovna, sv. 10) 5,250
copies printed.

Reviewer: Václav Tysl, Engineer; Ed.: Jan Nohejl, Doctor,
Engineer; Tech. Ed.: Marie Králová; Managing Ed. for Literature
on Electrical Engineering: František Káspár, Engineer, Doctor.

PURPOSE: The book is intended for readers with a secondary technical education.

COVERAGE: The author discusses the propagation of electromagnetic waves and describes antennas, waveguides, and microwave oscillators and amplifiers. He also discusses microwave spectroscopy and accelerators of elementary particles and explains the application of microwave techniques in medicine and astronomy. The author thanks Doctor Káspár for reviewing the text. There are 25 references of which 14 are Czech (including 5 translations), 4 English, 4 German and 3 Soviet.

Card 1/5

Czech/1562

Microwaves

TABLE OF CONTENTS:

Foreword	5
I. Electromagnetic Waves	13
1. Polarization of electromagnetic waves	14
2. Classification of electromagnetic waves according to frequency	16
II. Transmission of Microwaves by Means of Transmission Lines	18
3. Effect of load on input impedance of a transmission line	24
4. Attenuation of microwaves in transmission lines	32
5. Parallel-wire transmission line	35
6. Coaxial transmission line	43
7. General characteristics of waveguides	46
8. Electromagnetic field of various types of waves in waveguides	53
9. Practical application of waveguides	61
10. Strip-type waveguides	77
11. Cylindrical resonators	81

Card 2/5

Microwaves

Czech/1562

III. Propagation of Microwaves in Free Space	85
12. Electromagnetic horizon	86
13. Absorption of waves in the lower layer of the atmosphere	88
14. Reflection of microwaves in the ionosphere	91
15. Propagation of electromagnetic waves around the earth	97
IV. Oscillators and Amplifiers	99
16. Triode oscillators	100
17. Oscillators with modulation of electron velocity (klystrons)	104
18. Traveling-wave tubes	121
19. Diode magnetron oscillator	124
20. Multicavity magnetron	128
V. Antennas	144
21. General characteristics of directional antennas	144
22. Half-wave dipole	150
23. Dipole systems	157

Card 3/5

Microwaves

Czech/1562

24. Parabolic antennas (reflectors and radiators)	164
25. Antenna lenses	174
26. Other types of directional antennas	180
27. Turnstile antennas	185
VI. Indication and Measurement of Microwaves	187
28. Crystal diodes and their application to microwave technique	187
29. Receivers	189
30. Attenuation components	202
31. Standing-wave detector	209
32. Power measurement	212
33. Frequency measurement	216
34. Determination of antenna radiation pattern	219
VII. Some Instruments and Circuits Used with Microwaves	224
35. Radio communication	224

Card 4/5

Microwaves

Czech/1562

- | | | |
|-----|--|-----|
| 36. | Application of microwave techniques in medicine | 227 |
| 37. | Radar equipment | 229 |
| 38. | Application of microwave techniques in astronomy | 234 |

VIII. New Discoveries in Microwave Technique and Their Application in Modern Physics

- | | | |
|-----|---|-----|
| 39. | Microwave spectroscopy | 243 |
| 40. | Spectral lines | 244 |
| 41. | Experimental technique in radio spectroscopy | 271 |
| 42. | MASER generators | 275 |
| 43. | Some application of spectroscopy at microwave frequencies | 288 |
| 44. | Faraday's law as applied to centimetric waves | 295 |
| 45. | High-frequency accelerators of elementary particles | 300 |

Bibliography

Index

AVAILABLE: Library of Congress

JP/lrb
6-8-59

Card 5/5

HERDA, M., inz. CSc.; CESAK, K., inz.; WEBER, B., inz.; VYHNANEK, V., inz.;
KUNICKY, L., inz.; SIMEK, J., inz.; PROSTREDNIK, K., inz.

Maps for area planning and records of the built constructions.
Geod kart obzor 10 no.9/10:232-235 0 '64

WEBER, Bela

New plant fossil Upper Carboniferous gravels from the
Helvetian strata of the western part of the Mecsek Mountains.
Foldt kozl 94 no.3:379-381 J1-S '64.

WEBER, Bela

Thorium and rare earth indications in the Buda Mountains. Foldt
kezl 92 no.4:455-457 H-D '62.

WEDER D

Change of moisture content of drug seeds and fruits. *Datura Wimmeri* Mayer
Gyógyszerintézet. Társaság Esterházy 7, 194-7(1931).—(1) Seeds or fruits (about 50 cc.)
were dried above concd. H₂SO₄ and weighed. (2) The same were weighed again after
standing in air for several days (the air contained 23% moisture). (3) After 6 days
in a 24° thermostat in 100% moist air they were again weighed. The difference in the
values from (1) and (2) gives the max., that from (1) and (3) the max. content of moisture.
The normal moisture content averaged 4-5%, with a max. of 14-15%. *Ricinus com.*
memis seeds had 3 and 9% moisture contents, resp., *Atropa belladonna* 5 and 16%.
Archangelica officinalis 6 and 30%. etc. S. S. DE FINALLY

17

CA
WEBER, D.

Abstracts. Deutsches Institut für Angewandte Chemie.
Taschenrechner: 9, 307-30 (1963).—Abstracts made by
double annotation showed higher contents but were not
so perfect as those made by percolation. S.S. de F.

ASM-5LA METALLURGICAL LITERATURE CLASSIFICATION

130001 130002 130003 130004 130005 130006 130007 130008 130009 130010 130011 130012 130013 130014 130015 130016 130017 130018 130019 130020 130021 130022 130023 130024 130025 130026 130027 130028 130029 130030 130031 130032 130033 130034 130035 130036 130037 130038 130039 130040 130041 130042 130043 130044 130045 130046 130047 130048 130049 130050 130051 130052 130053 130054 130055 130056 130057 130058 130059 130060 130061 130062 130063 130064 130065 130066 130067 130068 130069 130070 130071 130072 130073 130074 130075 130076 130077 130078 130079 130080 130081 130082 130083 130084 130085 130086 130087 130088 130089 130090 130091 130092 130093 130094 130095 130096 130097 130098 130099 130100 130101 130102 130103 130104 130105 130106 130107 130108 130109 130110 130111 130112 130113 130114 130115 130116 130117 130118 130119 130120 130121 130122 130123 130124 130125 130126 130127 130128 130129 130130 130131 130132 130133 130134 130135 130136 130137 130138 130139 130140 130141 130142 130143 130144 130145 130146 130147 130148 130149 130150 130151 130152 130153 130154 130155 130156 130157 130158 130159 130160 130161 130162 130163 130164 130165 130166 130167 130168 130169 130170 130171 130172 130173 130174 130175 130176 130177 130178 130179 130180 130181 130182 130183 130184 130185 130186 130187 130188 130189 130190 130191 130192 130193 130194 130195 130196 130197 130198 130199 130200 130201 130202 130203 130204 130205 130206 130207 130208 130209 130210 130211 130212 130213 130214 130215 130216 130217 130218 130219 130220 130221 130222 130223 130224 130225 130226 130227 130228 130229 130230 130231 130232 130233 130234 130235 130236 130237 130238 130239 130240 130241 130242 130243 130244 130245 130246 130247 130248 130249 130250 130251 130252 130253 130254 130255 130256 130257 130258 130259 130260 130261 130262 130263 130264 130265 130266 130267 130268 130269 130270 130271 130272 130273 130274 130275 130276 130277 130278 130279 130280 130281 130282 130283 130284 130285 130286 130287 130288 130289 130290 130291 130292 130293 130294 130295 130296 130297 130298 130299 130300 130301 130302 130303 130304 130305 130306 130307 130308 130309 130310 130311 130312 130313 130314 130315 130316 130317 130318 130319 130320 130321 130322 130323 130324 130325 130326 130327 130328 130329 130330 130331 130332 130333 130334 130335 130336 130337 130338 130339 130340 130341 130342 130343 130344 130345 130346 130347 130348 130349 130350 130351 130352 130353 130354 130355 130356 130357 130358 130359 130360 130361 130362 130363 130364 130365 130366 130367 130368 130369 130370 130371 130372 130373 130374 130375 130376 130377 130378 130379 130380 130381 130382 130383 130384 130385 130386 130387 130388 130389 130390 130391 130392 130393 130394 130395 130396 130397 130398 130399 130400 130401 130402 130403 130404 130405 130406 130407 130408 130409 130410 130411 130412 130413 130414 130415 130416 130417 130418 130419 130420 130421 130422 130423 130424 130425 130426 130427 130428 130429 130430 130431 130432 130433 130434 130435 130436 130437 130438 130439 130440 130441 130442 130443 130444 130445 130446 130447 130448 130449 130450 130451 130452 130453 130454 130455 130456 130457 130458 130459 130460 130461 130462 130463 130464 130465 130466 130467 130468 130469 130470 130471 130472 130473 130474 130475 130476 130477 130478 130479 130480 130481 130482 130483 130484 130485 130486 130487 130488 130489 130490 130491 130492 130493 130494 130495 130496 130497 130498 130499 130500 130501 130502 130503 130504 130505 130506 130507 130508 130509 130510 130511 130512 130513 130514 130515 130516 130517 130518 130519 130520 130521 130522 130523 130524 130525 130526 130527 130528 130529 130530 130531 130532 130533 130534 130535 130536 130537 130538 130539 130540 130541 130542 130543 130544 130545 130546 130547 130548 130549 130550 130551 130552 130553 130554 130555 130556 130557 130558 130559 130560 130561 130562 130563 130564 130565 130566 130567 130568 130569

1ST AND 2ND CODES										3RD AND 4TH CODES									
PROCESSES AND PROPERTIES INDEX																			
WEBER, D.										17									
<p>Extracts of belladonna, hyocyanus and nux vomica and sirup of ferrum iodide. Deas6 Weber. Ber. ungar. pharm. Ges. 13, 392-402 (1847). Methods of prepn. are discussed in detail. S. S. de Finály</p>																			
ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION										E-27-52-14-14-14									
ROOM DIVISION										ROOM NUMBER									
SUBJECT										SUBJECT									

WEEDER, D
CA

Work in the spectroscopy laboratory. Density and the dry residues of tinctures. *Dens. Weber, Ber. usgar. pharm. Ges.* 9, 273-4 (1943); *Chem. Zentr.* 1943, II, 1980. —The sp. gr. and dry residues of various tinctures are recorded. It is recommended that besides the sp. gr. or alc. no. also the most modern methods for detg. tinctures (e. g., quartz luminescence analysis) should be used. S. Morgulis

17

ASTM, S. L. A. METALLURGICAL LITERATURE CLASSIFICATION

FROM SYMBOLS

INDEXED BY

CELLS

CELLS

NAJMAN, Emil, Dr.; WEBER, Danica, Dr.

Amebiasis and lambliasis in children. Lijec vjes 82 no.9/10:725-734
'60.

1. Iz Dječjeg odjela Opće bolnice "Dra M.Stojanovica" u Zagrebu
(AMEBIASIS in inf & child)
(GIRADIASIS in inf & child)

Webes, F.

Distr: 4E2c(f)

✓ Technical dyeing of polyamide and polyacrylonitrile textiles.
F. Weber (Prakt. Chem. 1931, 8, 192). — A review with comments on the streaky dyeings obtained when dyeing polyamide textiles with 2 : 1 Cr (or Co) complex dyestuffs, as well as with acid dyestuffs applicable in a neutral or weakly acid bath. Whereas it is possible to cover up unevenness on polyamide textiles dyed with 2 : 1 metal complex dyestuffs or with Alizarin Milling dyestuffs, by means of disperse dyestuffs, fastness must be taken into account. There are few really suitable vat dyestuffs available for this application. In the case of disperse (acetate rayon) dyestuffs, particle size is of great importance: the higher the degree of dispersion, the better the result. The limits of affinity between cation-active and the classical basic dyestuffs in the dyeing of polyacrylonitrile fibres is discussed. A table of the half-dyeing periods is given.

W. H. Kemp.

g-f

4
2 May
1

1/1

SCHMUTZER, E.; WEBER, G.

On the nonrelativistic second approximation of the Dirac theory of electron. Godishnik fiz mat 55 no.2:105-115 '60/'61 [publ. '62].

WEBER, G. 1949

(A Budapesti Egyetemi Korelettani Intezs-tebol)

"Lipolytic Power of the Serum in Laboratory Animals. Graphic Representation of the Monomolecular Velocity Constant of Lipase."

Orvosi Hetilap, 1949, 90/16(495-500)

Abst: Exc. Med. 11, Vol. 111, No. 4, p. 453

WEBER G. and DRECHSLER K. A Budapesti Egyetemi Korelettani Intezetabol. A laboratoriumi allatok lipolytikus kepessegerol. A lipase monomolekularis sebessegi allandojanak grafikus meghatarozasa, Lipolytic power of the serum in laboratory animals. Graphic representation of the monomolecular velocity constant of lipase, Orvosi Hetilap, Budapest 1949, 90/16 (495-500) Graphs 4 Tables 1

The resistance of laboratory animals against tubercle bacilli does not run parallel to the level of serum lipase. The white rat, which is very resistant to tubercle bacilli, shows the lowest level of serum lipase, whereas the rabbit, which is very sensitive to the bovine type, and the guinea-pig, which is sensitive to the human type, have respectively two and three times as much serum lipase as the white rat. A rapid and easy graphic method of estimating the monomolecular velocity constant of lipase is given. The stalagmometric estimation of the lipase is reduced to 22 minutes (formerly many hours) and the very tedious procedure is simplified.

Olbrich - Edinburgh (II, 4, 6, 15)

So: Medical Microbiology and Hygiene, Section IV, Vol 3, No 1-6

LENKEI, Peter; WEBER, Gyorgy

Some interesting aspects in the design and construction of a
storage building. Magyar Építész Lapok 13 no.12:743-748 '64.

L 29456-66 IJP(c) JT

ACC NR: AP5028059

SOURCE CODE: GE/0030/65/012/001/0071/0080

AUTHOR: Egorov, W. D., ~~Lomonosov University, Moscow (Lomonosov-Universität Moskau)~~; Muller, G.O.; Weber, H. 53
B

ORG: ~~The~~ Physical-Technical Institute, German Academy of Sciences, Berlin
Division of Crystal Electronics (Physikalisch-Technisches Institut
der Deutschen Akademie der Wissenschaften, Abteilung
Kristallelektronik); Lomonosov University, Moscow (Lomonosov-Universität)
[Egorov]

TITLE: Cathodoluminescence of CdS under the effect of strong cathode ray excitation

SOURCE: Physica status solidi, v. 12, no.1, 1965, 71-80

TOPIC TAGS: crystal, crystal impurity, cathode ray, electron beam, current density, emission spectrum, ELECTRON ENERGY, CATHODOLUMINESCENCE

ABSTRACT: The purpose of the experiments described here was to verify the occurrence of certain effects on Frerichs-crystals with natural faces at electron energies under 10 keV. High purity was to be expected from the crystals grown by a modified Frerichs method through the reaction between Cd vapor and H₂S, and low energy electron bombardment of constant beam current density gave high excitation density

Card 1/2

L 29456-66

ACC NR: AP5028059

because the depth of penetration increases quite linearly with the electron energy. An attempt was made to repeat Basov's experiments to stimulate emission from CdS crystals at 90° K by cathode ray bombardment. No stimulated emission was observed. The intensity of blue emission increases with the electron energy and saturation of the green emission occurs in the 21. to 8.5ke V electron energy range, but at different energies for each crystal. The blue emission prevails at high excitation densities for all electron energies, but the structure of the green emission band remains essentially constant. The experimental results suggest a relation between the green emission and the surface properties of the crystals. "We thank the Ministry of University Affairs of the USSR (Ministerium fur Hochschulwesen der UdSSR) and the Ministry of National Education of the GDR (Ministerium fur Volksbildung der DDR) for making it possible for one of us, through a six month leave of absence, to participate in this study at PT (The Physical Technical Institute). Orig. art. has: 8 figures.

SUB CODE: 20/ SUMB DATE: 29Jul65/ ~~SECRET~~ / SOV REF: 003/ OTH REF: 010

Card 2/2

COUNTRY : Czechoslovakia H-35
 CATEGORY :
 ABS. JOUR. : RZKhim., No. 1959, No. 88875
 AUTHOR : Weber, H.
 INST. :
 TITLE : Some Czechoslovak Synthetic Tanning Agents
 from Dihydric Phenols
 ORIG. PUB. : Kozarstvi, 1958, 8, No 8, 238-239
 ABSTRACT : Description of the production of synthetic
 tanning agents from by-products obtained in processing of
 coal, primarily diphen and optol. Diphen or optol are a
 mixture of pyrocatechol, methyl-pyrocatechol and isomethyl
 pyrocatechol. The pyrocatechols are the basic raw material
 for the production of syntans. The tanning agent Kortan Q1
 obtained by processing of pyrocatechols has been tested
 with favorable results at the leather factories of Czecho-
 slovakia and USSR, where it is known as Syntan No 4. In
 recent years there have been obtained syntans of the Kortan
 type with a minimal content of sulfo-groups and a maximum
 content of dihydric phenols. They produce greater tanning
 CARD: 1/2

295

COUNTRY : Czechoslovakia
CATEGORY :

H-35

ABS. JOUR. : RZKhim., No. 1959, No. 88875

AUTHOR :
INST. :
TITLE :

ORIG. PUB. :

ABSTRACT : and filling effect. Their price does not exceed the price of imported natural tanning agents. In making light-colored leather pyrosulfite is added to the liquid Kortan. It is possible further to improve the properties of syntans of this type. -- M. Lyuksemburg.

CARD: 2/2

L 29661-65 EWT(1)/EWT(2)/T/EWP(1)/EWP(2) LJP(1) JD

ACCESSION NR: AP5001295

P/0045/64/026/03-/0737/0744

AUTHOR: Muller, G. O.; Weber, H.

TITLE: Effect of electric fields on the cathodoluminescence of CdS single crystals

SOURCE: Acta physica polonica, v. 26, no. 3-4, 1964, 737-741

TOPIC TAGS: cathodoluminescence, electric field enhanced luminescence, cadmium sulfide single crystal, metal contact, electrode, electron beam, electric field effect, luminescence, field polarity, quenching, barrier layer, field influenced recombination

ABSTRACT: The cathodoluminescence of pure undoped CdS single crystals with metal contacts in a sandwich arrangement under the influence of weak d-c fields with electron beams of 5--25 kev impinging on one electrode were studied in order to determine the effect of the electric fields and the relationship between d-c field effects, electrical behavior, and type of contact on the irradiated surfaces. One electrode in the measuring apparatus always consisted of indium and was always ohmic, while the irradiated electrodes were of Al, Au, Ti, Cr, or Te. It has been found that all crystals with measurable field effect have a short-circuit current

Card 1/2

L 29661-65

ACCESSION NR: AP5001296

4

during bombardment with electrons, while crystals without a short-circuit current do not display any measurable field effect at applied voltages of up to 50 v. Field-enhanced luminescence and ~~un~~ dependent only on the polarity of the field are observed for the same crystal with the same cathode excitation. Field factors showing the magnitude and type of effect the electric field has on cathodoluminescence (determined as the ratio of intensity with field to intensity without field) in the 0.2 - 4 range for average fields below 10^4 v/cm are common. The relationships between the field factors and applied voltages differ with the spectral emission bands. An interpretation of the measurements is suggested by the qualitatively different behavior of the crystals assuming a field-influenced recombination in the barrier layer. "The authors thank Dr. F. Eckart and Dr. E. Schurer for their many valuable discussions, and Dr. L. Hildisch for preparing the crystals." Orig. art. has: 5 figures.

ASSOCIATION: Institute of Technical Physics, German Academy of Sciences, Berlin, German Democratic Republic

SUBMITTED: 00

ENCL: 00

SUB CODE: 55, OP

NO REF SOV: 000

OTHER: 006

Card 2/2

WEBER, H. dr. (Csehszlovakia)

Solvent tannage in the Czechoslovak Socialist Republic. Ber Cipe 13
no.6:179-181 N. '63.

NICOLAU, Edm.; WEBER, I.; GAVAT, St.

Apparatus for automatic recognition of the vowels. Automatica
electronica 7 no.6:255-261 N-D'63.

WEBER, Ilja, MUDr

Cancer of the cervix uteri in Soviet literature. Lek. listy 9
no.14:332-333 JI '54.

1. Z II. Porodnicko gynekologicke kliniky, prednosta prof.
Dr Horalek.
(CERVIX, UTERINE, neoplasms,
*review, Russia)

Weber, I.

3

LAPIDAT TEOPTIL, NARAI.
Bucuresti (in copy); Given Name

Country: Rumania

Academic Degree: Dr.

Affiliation: Veterinary Hospital (Spitalul Veterinar), Tg. Secuiesc.

Source: Bucharest, Probleme Zootehnice si Veterinare, No 4, 1961,
pp 60-61.

Date: "The Treatment of a Fetal Enostic of Neurobacillosis in
Bulls."

Co-authors:

WEBER, I., Dr., The People's Council of the Raion of Tg. Secuiesc
(Sfatul Popular al Raionului, Tg. Secuiesc).

WEBER, Istvan

Experiences gained in processing fur materials. Bor cipo
10 no.2:50-52 Mr '60.

1. Szegedi Szorcs- es Borruhakeszito Vallalat.

WEBER, J.

"Remarks on the Article 'Problems of Producing Lumber Products of Good Quality'", P. 166, (FAIPAR, Vol. 4, No. 6, June 1954, Budapest, Hungary)

SO: Monthly List of East European Accessions (EEAL), LC, Vol. 4, No. 3, March 1955, Uncl.

HORVATH, Z., doctor of eng.sc.; WEBER, J.

The Parkes process with zinc addition in two stages granting minimum zinc consumption. Acta techn Hung 40 no.3/4:263-284, '62.

1. Metallurgical Department of the University for Heavy Industry, Miskolc.

WEBER, J. ; KOUTECKY, J.

Kinetics of electrode processis. XV. Tables for calculating polarographic currents due to depolarization processes involving preceding or consecutive rapid monomolecular chemical reactions. p. 562

CHEMICKÉ LISTY (Ceskoslovenska akademie ved. Ceskoslovenska spolecnost chemicka) Praha, Czechoslovakia. Vol. 49, no. 4, Apr. 1955

Monthly List of East European Accessions (EEAI) IC. Vol. 9, no. 1/1960
Uncl.

VELEK J.

COUNTRY	:	Czechoslovakia	B-12
CATEGORY	:	Physical Chemistry--Electrochemistry.	
ABS. JOUR.	:	RZKhim., No. 14 1959, No.	48900
AUTHOR	:	<u>Veber, J.</u> ; Kuta, J. and Smoler, I.	
INST.	:	Not given	
TITLE	:	Instantaneous Polarographic Currents. I. Diffusion-Limited Currents and Currents Limited by Slow Electrode Reactions. II. Investiga*	
ORIG. PUB.	:	Chem Listy, 51, No 7, 1249-1258; 1259-1263 (1958)	
ABSTRACT	:	I. Instantaneous polarographic currents and their time dependence are more suitable for use in the quantitative analysis of the kinetics of electrode processes than the average polarographic currents recorded by the classic polarographs. The author discusses a number of expressions which can be used for the representation of some types of instantaneous polarographic currents, taking into account spherical diffusion (SD). For a reversible redox system in	
CARD: 1/9	:	* tion of Current-Time Curves at Various Half-Wave Potentials for Reversible and Irreversible Processes.	

COUNTRY : Czechoslovakia
CATEGORY :

B-1

ABS. JOUR. : RZKhim., No. 14 1959, No.

48900

AUTHOR :
INST. :
TITLE :

ORIG. PUB. :

ABSTRACT : which the product of the electrode reaction dif-
fuses back into the solution, the instantaneous
current is higher than the current calculated
by the Ilkovic equation at all values of the
potential. Following correction for SD, the
dependence of the instantaneous current on the
time over the interval 1-6 sec is expressed by
the relation

$$i = k \cdot t^{0.192}$$

In the case of amalgam formation the current

CARD: 2/9

13-57

COUNTRY : Czechoslovakia B-12
CATEGORY :
ABS. JOUR. : AZKhim., No. 14 1959, No. 48900
AUTHOR :
INST. :
TITLE :
ORIG. PUB. :
ABSTRACT : after the application of the SD correction is increased when the potential is more negative than the normal potential and is decreased when the potential is more positive than the normal potential. This difference is due to the difference in the diffusion spaces of the reduced and the oxidized forms. If the electrode process proceeds slowly, the SD correction can be neglected in the first approximation; the character of the dependence of the depolarization current

CARD: 3/9

COUNTRY : Czechoslovakia
CATEGORY :

B-12

ABS. JOUR. : RZhKhim., No. 14 1959, No.

48900

AUTHOR :
INST. :
TITLE :

ORIG. PUB. :

ABSTRACT : on the time depends on the potential; the slope of the $\log i - \log t$ curve lies in the range $1/6 - 2/3$. At lower overpotentials the above slope may attain a maximum value in the region of the half-wave potential.
II. Since the effect of concentration polarization on the first and on subsequent drops is unequal (RZhKhim, No 16, 1954, 37501; No 21, 1954, 46177; No 4, 1956, 9475), the current-time curves for the first drops, which are unaffected by

CARD: 4/9

3-60

COUNTRY	:	Czechoslovakia	B-12
CATEGORY	:		
ABS. JOUR.	:	RZKhim., No. 14 1959, No.	48900
AUTHOR	:		
INST.	:		
TITLE	:		
ORIG. PUB.	:		
ABSTRACT	:	<p>preceding polarization, must be recorded. The recording of these curves must be carried out in solutions containing a depolarizer in concentrations of $(3-5) \cdot 10^{-3}$ M at potentials corresponding to the following $i/i(\text{lim})$ ratios: 0.75, 0.5, 0.25, and 0.1. For reversible systems, e.g., Fe(II) — Fe(III) and Ti(III)-Ti(IV), in the presence of oxalate, and quinone-hydroquinone in phosphate buffer at pH 6.8, the authors have found a linear dependence between</p>	

CARD: 5/9

COUNTRY : Czechoslovakia
 CATEGORY : 8-12
 ABS. JCUR. : RZKhim., No. 14 1959, No. 48900
 AUTHOR :
 INST. :
 TITLE :
 ORIG. PUB. :
 ABSTRACT : log i and log t. In agreement with theory, the slopes of these curves are 0.195 (independent of E). For reversible electrode processes in which the reduced form forms an amalgam, the linear relationship between log i and log t was found to be still valid, but the slope was found to correspond to the value cited only for $i(\lim)$; at more positive values of E the slope decreases and at $E_{1/2}$ was found = 0.167 (zero SD correction). Under these conditions the current is

CARD: 6/9

B-61

COUNTRY	:	Czechoslovakia	B-12
CATEGORY	:		
ABS. JOUR.	:	RZKhim., No. 14 1959, No.	48900
AUTHOR	:		
INST.	:		
TITLE	:		
ORIG. PUB.	:		
ABSTRACT	:	<p>equal to that calculated by the Ilkovic equation. At even more positive values of E, the slope gradually decreases (negative SD correction). At such E, when $i < 0.1 i(\text{lim})$, the difference between the current-time curves for the first and the subsequent drops decreases. For irreversible electrode processes at high overvoltages the value of the parameter giving the parabolic dependence between i and t decreases uniformly from the foot of the wave</p>	

CARD: 7/9

COUNTRY	:	Czechoslovakia	B-12
CATEGORY	:		
ABS. JOUR.	:	RZKhim., No. 14 1959, No.	48900
AUTHOR	:		
INST.	:		
TITLE	:		
ORIG. PUB.	:		
ABSTRACT	:	<p>where it is $2/3$) with increasing i and attains a value of $1/5.3$ at $i(\lim)$. The dependence between $\log i$ and $\log t$ is linear only for values of i which are small compared to $i(\lim)$, in agreement with theory. At the foot of the wave the current is affected by the rate of the electrode reaction and at more negative E the rate of diffusion also exerts an effect. For irreversible electrode processes at low overpotentials, the value of the slope passes</p>	

CARD: 8/9

2-62

COUNTRY : Czechoslovakia B-12
CATEGORY :
ABS. JOUR. : RZKhim., No. 14 1959, No. 48900
AUTHOR :
INST. :
TITLE :
ORIG. PUB. :
ABSTRACT : through a maximum as i increases. Examples discussed are the reduction of $Zn(2+)$ against a background of Na_2SO_4 and the reduction of $Eu(3+)$ against a background of $NaClO_4$ and $NaCl$. Metals of the Fe group exhibit anomalous current-time curves.

P. Zuman

CARD: 9/9

AUTHOR: Weber, Jan

C Z/8-52(82)-10-8/39

TITLE: Polarographic Currents in the Case of Periodically Alternating Voltages (Polarografické proudy při periodicky proměnném napětí). Part II. Theory of Currents in the Case of Regeneration of the Depolariser by Chemical Reaction in the Solution (II. Teorie proudů při regeneraci depolarisátoru chemickou reakcí v roztoku)

PERIODICAL: Chemické Listy, 1958, Vol 52(82), Nr 10, pp 1888-1898 (Czechosl)

ABSTRACT: This is a continuation of work published by another author, J. Koutecký (Chem. listy, Vol 49, p 1454, 1955: Collection, Vol 21, p 433, 1956). Calculation of the current fed into a perfectly electro-active system of redox, the oxidised and reduced components of which pass into solution as a result of a pseudo-monomolecular reaction, is one of the simplest problems in the theory of polarographic currents in the case of applying a voltage which is variable with time. Its solution provides a possibility of accurate calculation of the kinetic currents, even in the case of an applied voltage which varies periodically, and could be used as a theoretical basis for experimental investigation of chemical reactions which lead to repeated regeneration of the depolariser. Polarisation by a periodically varying voltage permits investigating the entire system under experimental conditions which differ from those pertaining to polarography with a constant voltage and it may also contribute to elucidating the complicated chemical reactions involved in these

Card 1/2

Z/8-52(82)-10-8/39

Polarographic Currents in the Case of Periodically Alternating Voltages.
Part II. Theory of Currents in the Case of Regeneration of the Depolariser by Chemical Reaction in the Solution

processes. In this paper, formulae are derived for the instantaneous current corresponding to the electrode reaction in the case that the reaction product is regenerated to the original depolariser by a chemical reaction in the solution. The calculations are made for depolarisation currents of an ideally reversible redox system for plane stationary spherical and increasing drop electrodes with any time characteristic of the applied potential. The general formulae were specially adapted for a potential with a periodically square-topped shape and the characteristic of the dependence of the instantaneous current on time for some simpler cases is calculated. The instantaneous and the average current after establishment of the stationary state are calculated. Acknowledgments are made to Dr. J. Koutecký for his interest in the work and M. Čejková for carrying out some of the numerical calculations. There are 2 figures and 1 table and 11 references, 4 of which are Czech, 5 English, 2 German.

ASSOCIATION: Ústav fyzikální chemie, Československá akademie věd, Praha
(Institute of Physical Chemistry, Czechsl. Ac.Sc., Prague)

Card 2/2

COUNTRY	:	Czechoslovakia	B-12
CATEGORY	:		
ABS. JOUR.	:	RZKhim., No. 5 1950, No.	17167
AUTHOR	:	Weber, J.	
INST.	:	Not given	
TITLE	:	Polarographic Currents with Periodically Varying Potentials. I. Theoretical Relations for the Current During the Regeneration of the Depolarizer	
ORIG. PUB.	:	Collection Czechoslov Chem Commun, No 6, 1770-1782 (1959)	
ABSTRACT	:	See RZhKhim, 1959, No 25, 81546	

CARD: 1/1 * by Chemical Reactions₅₅ in the Solution.

SOURCE	: Czechoslovakia	B-12
CATEGORY	:	
ABS. JOUR.	: RZhKhim., No. 5 1960, No.	17166
AUTHOR	: Weber, J. (I); Kuta, J. and Smoler, I. (II)	
INSTR.	: Not given	
TITLE	: Instantaneous Polarographic Currents. I. Diffusion-Limited Currents and Currents Limited by Slow Electrode Reactions. II. Investigation of	
ORIG. PUB.	: Czechoslov Chem Commun, 24, No 5, 1424-1435; No 7, 2208-2220 (1959)	
ABSTRACT	: See RZhKhim, 1959, No 14, 48900.	

CARD: 1/1 *the Curves Expressing the Dependence of the Current on the Time at Various Potentials of the Polarographic Wave for Reversible and Irreversible Processes.

WEER, J.

"Instantaneous polarographic currents." I. Currents governed by diffusion and currents governed by slow-electrode reaction. In German. p. 1424.

COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS, Praha, Czech.
Vol. 24, No. 5, May 1959

Monthly List of East European Accessions (EEAI), LC, Vol. 8, No. 6, Sept, 59

Unclassified

WEBER, J.

Polarographic currents at periodically varying potential. II.
Calculation of stationary currents with rectangular pattern of the
polarizing potential. In German. Coll.Cz.Chem. 24 no.9:3041-3045
S '59. (EEAI 9:5)

1. Institut für physikalische Chemie, Tschechoslovakische Akademie
der Wissenschaften, Prag.
(Polarograph and polarography) (Electric currents)

KOUTECKY, J.; WEBER, J.

The theory of polarographic currents affected by the adsorption of electro-inactive substances in a reversible electrode reaction. Coll Cz Chem 25 no.5:1123-1126 My '60.

1. Institut für physikalische Chemie, Tschechoslowakische Akademie der Wissenschaften, Prag.

WEBER, J.; KOUTECKY, J.

Effect of absorbable substances on polarographic currents. V. Instantaneous and delayed currents in the presence of charged and uncharged surface-active substances undergoing slow adsorption. Coll Cz Chem 25 no.12:2993-3003 D '60. (EEAI 10:9)

1. Institut für physikalische Chemie, Tschechoslowakische Akademie der Wissenschaften, Prag.

(Polarograph and polarography) (Surface-active substances)
(Adsorption)

VAGUNDA, Simon; REZAC, Milos; WEHER, Jaromir

Gastric cancer in a 12-year-old boy. Cesk.pediat.15 no.10:913-916
0'60.

1. Patologicko-anatomicke oddeleni, prednosta prim.dr. S.Vagunda
Centralni rtg oddeleni, prednosta prim.dr. E. Sanda. Chirurgicke
oddeleni, prednosta prim.dr. Fr. Kudlac.
(STOMACH NEOPLASMS in inf & child)

L 43542-65 EWG(v)/EWT(1) Pe-5/Po-4/Pq-4/Pac-4/Pae-2 GN/GS
UR/0000/63/000/000/0132/0142

ACCESSION NR: AT5009182

AUTHOR: Zverev, M.S. (Corresponding member AN SSSR); Hemiro, A.A.

TITLE: Observation program of the Soviet astrometric expedition to the southern hemisphere

SOURCE: Astrometricheskaya konferentsiya SSSR. 15th. Pulkovo, 1960. Trudy. Moscow, Izd-vo AN SSSR, 1963, 132-142

TOPIC TAGS: astrometry, meridian observation, astrograph, star catalogue, fundamental star, reference star, astronomical instrument

ABSTRACT: The problems involved in compiling a Catalogue of Faint Stars are discussed in detail, particularly the necessity of making the necessary observations in the southern hemisphere. The work already done by southern hemisphere observatories is discussed, as are the repeated resolutions passed at successive International Astronomical Union congresses, often on the recommendation of Soviet astronomers, calling for an expansion of such observations (the principles of compilation of such a catalogue require observations of each star, galaxy and minor planet at not less than 3 observatories). As a result of an agreement reached between the Soviet Academy of Sciences and the Chilean government after a visit of the Rector of Santiago University to the Soviet Union in the

Card 1/3

L 43542-65

ACCESSION NR: AT5009182

spring of 1960, an astrometric expedition is to be sent to Cerro Calan Observatory in Chile; specialists from the Pulkovskaya Observatoriya (Pulkovo Observatory) will work there in close collaboration with Chilean astrometrists. The following work will be done in the field of meridian astrometry. 1. Absolute determination of the right ascensions of bright and faint fundamental stars using a large transit instrument. 2. Absolute determination of the declinations of these same stars using a vertical circle. 3. Observations of the sun and major planets, especially Mercury, Venus and Mars, using a large transit instrument and a meridian circle. 4. Differential connection of the system of right ascensions of stars in the northern and southern sky using a transit instrument with a broken telescope. 5. Differential determination of the coordinates of faint and bright reference stars using a meridian circle. The methods to be used in implementing this program are discussed. Particular attention is given to possible methods which can be used for absolute determination of the azimuth of the large transit instrument which is to be employed. The meridian circle will be used in observing about 11,500 faint reference stars and 2,000 bright stars in the zone -25° to $+90^{\circ}$. The meridian instrument available at Cerro Calan is a Repsold instrument with an objective diameter of 19 cm, a circle 75 cm in diameter and with $4'$ graduations. Plans call for a double-meniscus (Maksutov) astrophotograph in making photographic observations of southern galaxies. Orig. art. has: 1 table.

Card 2/3 Submitted 06 Apr 63

ZVEREV, M.S.

Photographic method of determining the corrections to all the
diameters of a divided circle. Astron. zhur. 41 no.6:1128-1137
N-D '64 (MIRA 18:1)

1. Glavnaya astronomicheskaya observatoriya AN SSSR.

S WEBER, J.

18

The Use of High-Quality Cast Iron as a Constructional Material. J. Weber. (Practical Engineering, 1961, 1, Sept., 241-242). [LIT-P0001]. The author directs attention to the necessity for more research into the properties of high-quality cast iron and for publishing standards to enable engineers to use this material more widely.--V. O.

WEBER, J.

Computation of the activity of water and of the activity coefficient in the Na OH-H₂O-system. p. 564.

(KOHASZATI LAPOK, Budapest, Vol. 9, no. 12, Dec. 1954.)

SO: Monthly list of East European Accessions, (EEAL), LC, Vol. 4, no. 1, Jan. 1955,
Uncl.

WEBER, Jozsef, okleveles femkohomernok

Calculation of the activity and activity coefficient of water
in the $\text{NaOH-H}_2\text{O}$ system. Kohlap 9 no. 12: 564-576 D '54.

1: Femkohaszati Tanszek.

WEBER, J.

Distr: 4E2c

6483

621.784.422.5

* Weber J. Phosphating Steel.

"Fosforanowanie stali" Warszawa, 1938, PWT, 16°, 184 pp., 102 figs., 41 tabs.

Theoretical fundamentals and technology of the process of phosphating steel. Desirable equipment and conditions of operation are given.

EW
11

OM

2
1
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

WEBER, J.

Conditions of equilibrium of the magnesium oxide system. p.174. (Mohaszati Lapok. Budapest, Vol. 11, no. 4, Apr. 1956.)

SO: Monthly List of East European Accessions (EEAL) LC., Vol. 6, no. 7, July 1957 Uncl.

WEBER, J.

HUNGARY / Physical Chemistry. Thermodynamics.
Thermochemistry. Equilibrium. Physical
Chemical Analysis. Phase Transitions.

B

Abs Jour: Ref Zhur-Khimiya, No 19, 1958, 63663

Author : ~~Weber Jozsef~~

Inst : Not given

Title : Why the Temperature Coefficient of the Thermo-
dynamic Normal Potential of the Reaction of the
Formation of Carbon Monoxides Differs From
Coefficients Belonging to Reactions of Metal
Oxides.

Orig Pub: Kohasz. lapok, 1957, 12, No 3, 105-108

Abstract: A thermodynamic normal potential of the reaction

Card 1/5

HUNGARY / Physical Chemistry. Thermodynamics.
Thermochemistry. Equilibrium. Physical
Chemical Analysis. Phase Transitions.

B

Abs Jour: Ref Zhur-Khimiya, No 19, 1958, 63663

Abstract: $2C + O_2 = 2CO$ (I), in contrast to the reaction of the formation of metal oxides, will become more negative with an increase of temperature, from which it follows that the equilibrium constant K must increase with the temperature. However, this contradicts Le-Shatel'ye's principle, since the reaction is exothermic. The author explains this contradiction on the basis of an analysis of data concerning the equation for the dependence of molecular heat on the temperature under gradual pressure, for the enthalpy and entropy of the formation of reaction components under standard conditions, for molecular heat, for the enthalpy and entropy of reaction,

Card 2/5

HUNGARY / Physical Chemistry. Thermodynamics.
Thermochemistry. Equilibrium. Physical
Chemical Analysis. Phase Transitions.

B

Abs Jour: Ref Zhur-Khimiya, No 19, 1958, 63663

Abstract: as well as the dependence of the normal reaction potential on the temperature, and the dependence between the normal potential and $\lg K$. From the latter dependence, it is clear that K and $\lg K$ are very large numbers, but, with an increase of temperature, they decrease sharply. This indicates an increase of the partial pressure of O_2 due to the partial pressure of CO , in spite of the fact that the normal potential becomes more negative. The explanation of the contradiction consists of the following. The

Card 3/5

HUNGARY / Physical Chemistry. Thermodynamics.
Thermochemistry. Equilibrium. Physical
Chemical Analysis. Phase Transitions.

B

Abs Jour: Ref Zhur-Khimiya, No 19, 1958, 63663

Abstract: entropy of reaction (I), in contrast to the
entropy of the formation of oxides of metals, is
a large positive number. As a result of this, in
the equation $\Delta G_T^O = \Delta H_T - T \Delta S_T$ at every
temperature, and with an increase in T , ΔG_T^O
becomes more negative. But in addition to this,
the value K increases only under the condition
 $\Delta G_{T_2}^O - \Delta G_{T_1}^O > -4.575(T_2 - T_1)$ which, for the given
equation, is not achieved. But the increase of
the negative value ΔG_T^O lies in the fact that,
with an increase of temperature, the carbon is

Card 4/5

Card 5/5

6

WEBER, JOSEF

Distr: 4E3d 1

Temperature coefficient of the normal potential of carbon monoxide formation in relation to that of other monoxides.
 Josef Weber. *Kohászati Lapok* 90, 105-8 (1957).—The entropy of the reaction $2C + O_2 = 2CO$ is pos. and high, contrary to that of reactions resulting in the formation of metal oxides. The thermodynamic normal potential ($\Delta G_f^\circ = \Delta H_f^\circ - T\Delta S_f^\circ$, where ΔH_f° is the enthalpy and ΔS_f° is the entropy of the reaction) is at all temps. higher than ΔH_f° , and upon increasing the temp. ΔG_f° will become more neg. The relation between ΔG_f° and the equil. const. (K) of the reaction is $\Delta G_f^\circ = -RT \ln K = -4.575T \log K$, which indicates that K increases with increasing temp. only if $\Delta G_f^\circ - \Delta G_{CO}^\circ > -4.575(T_2 - T_1)$. This is not the case with the reaction resulting in CO formation; therefore, with increasing temp. C tends to be oxidized to CO rather than to CO_2 . The foregoing relation is utilized in *metal smelting*, as it permits the reduction of metal (M) oxides with C at appropriate temp., since above that temp. $\Delta G_f^\circ = \Delta G_{CO}^\circ - \Delta G_{MO}^\circ$ would be neg. and a spontaneous reaction would result.

L. G. Aryai

WEBER, J.

Distr: 4E2c

¹⁹ ²⁷ ^{16/54}
~~Reduction of metal oxides with carbon monoxide.~~
~~Webster, Kohászai Lapok 91, 122-30 (1958).—The relations~~
~~between the thermodynamical normal potential (I), CO/CO~~
~~equil. const. (II), and temp. were studied for various metal~~
~~oxides. The reduction of Fe₂O₃, Mn₂O₃, MoO₃, Na₂O,~~
²⁷ ~~NiO, PbO, TiO₂, V₂O₅, and WO₃ is exothermic, II decreasing~~
~~and I becoming more neg. with an increase in temp. The re-~~
~~verse conditions exist for Cr₂O₃, MnO, and VO. The equil.~~
~~is affected by temp. changes, and whether the reaction is~~
³⁸ ~~exothermic or endothermic, are not detd. by the direction~~
~~of the shift of I with an increase in temp. The effect of~~
^{1/} ~~temp. on II is detd. by the exothermic or endothermic na-~~
~~ture of the reaction.~~ L. G. Arvai

PM Jf

COUNTRY : POLAND H
 CATEGORY : Chemical Technology. Chemical Products and
 Their Applications. Corrosion. Corrosion*
 ABS. JOUR. : RZKhim., No. 23 1959, No. 82618
 AUTHOR : Weber, J.
 INST. : -
 TITLE : Study of Non-Scale Forming Solution for Phos-
 phatizing.
 ORIG. PUB. : Prace Inst. mech., 1959, 7, No 25, 69-76
 ABSTRACT : Addition of $H_2C_2O_4$ and its salts into a phos-
 phatizing (P) bath tends to prevent forma-
 tion of scale in the P process. Thus obtained
 films (PF) do not affect mechanical properties
 of lacquer layers applied over them. The
 quality of FF deteriorates with the increased
 concentration of $H_2C_2O_4$ and the promoter
 ($NaNO_3$, $NaNO_2$, $NaClO_3$). The non-scaling so-
 lutions may be employed for P of Zn-Al alloys.

*Control.

CARD:

H - 16

COUNTRY : H
CATEGORY :
ABS. JOUR. : RZKhim., No. 23 1959, No. 82618
AUTHOR :
INST. :
TITLE :
ORIG. PUB. :
ABSTRACT : Such solutions are also applicable for P of
Con'd Al. Corrosion resistance of PF obtained from
non scaling solutions, particularly at ele-
vated temperatures (500), is lower than of PF,
obtained from common baths. -- V. Levinson.

CARD: 2/2

COUNTRY : POLAND H
 CATEGORY : Chemical Technology. Chemical Products and
 Their Applications. Corrosion. Corrosion *
 ABS. JOUR. : RZKhim., No. 23 1959, No.82628
 AUTHOR : Biestek, T.; Web er, J.
 INST. : -
 TITLE : Accelerated Corrosion Tests of Phosphate
 Films on Steel
 ORIG. PUB. : Prace Inst. mech., 1959, 7, No 25, 77-85
 ABSTRACT : The results of tests conducted with phosphste
 films (PF) on steel, obtained from baths of
 different composition indicate that the cor-
 rosion resistance (CR) of PF, covered with
 protective lacquer layers, increases with in-
 creased PF thickness. Superior CR possess PF
 obtained from solutions of "Mazhef" or "Perkar
 2" salts ($ZnHPO_4 + H_3PO_4 + Cu$ salt). PF, obtai-
 ned from other, including the non-scale for-
 ming solutions, possess considerable lower CR.
 *Control.
 CARD: 1/2

COUNTRY :	H
CATEGORY :	
ABS. JOUR. :	RZKhim., No. 23 1959, No. 82628
AUTHOR :	
INST. :	
TITLE :	
ORIG. PUB. :	
ABSTRACT :	It is noted that thinner PF, for instance
Con'd	those obtained from non-scale forming solu- tions, have better mechanical properties, as compared to other PF, particularly the adhe- sive properties. The drop method proved use- ful only for testing PF of comparatively grea- ter thickness. Of the accelerated testing me- thods the most rugged and rapid is that per- formed with the use of a salt chamber, the least rugged - the test performed in humid atmosphere. -- V. Levinson.
CARD:	2/2
	H - 18

HORVATH, Zoltan, dr.; ~~WEBER, Jozsef~~

Thermodynamic examination of the processes occurring during the
dezincification of lead by means of chlorine gas. Koh lap 93 no. 5:
193-199 My '60.

21 3
✓ Thermodynamics of the dezincification of lead with
gaseous chlorine. II. Zoltán Horváth and József Weber.
Köhdzsalí Lapok 15, 268-73 (1960); cf. *CA* 54, 18264c.
From the equil. const. calcd. from the thermodynamic
normal potential, the changes in the equil. Zn content of the
refined Pb, in relation to the temp. and to the Pb content of
the chloride slag, can be calcd. An increase in the Pb con-
tent and (or) a decrease in temp. will reduce the Zn content
of the refined product. L. G. Arvai

HORVATH, Zoltan, dr.; WEBER, Jozsef

The Parkes process as a work method insuring double addition
of zinc in special view of the least use of zinc. Koh lap
95 no.3:108-114 Mr '62.

HORVATH, Zoltan, dr.; WEBER, Jozsef

Determination of the working method ensuring the lowest zinc consumption in the Parkes process with residual foam recycling and single zinc addition. Koh lap 96 no.5:216-221 My '63.

1. Magyar Tudományos Akademia Kohaszati Munkakozossege.